CLIMATOLOGICAL TABLE—Summary of the Records of Temperature, Rainfall, and Sunshine for August 1971—continued

Station	Height of Station Above M.S.L.	Air Temperatures in Degrees (Celsius)								Rainfall in Millimetres					
		Means of		Mean of A	Differ-	Absolute Maximum and Minimum			Total	No.	Differ-	Maximum Fall		Bright Sun- shine	
		A Max.	B Min.	and B	From Normal	Maxi- mum	Date	Mini- mum	Date	Fall	Rain Days	From	Amount	Date	
	Metres	°C	°c	°c		°c		°c		mm			mm		Hrs
Taieri Berwick Forest Dunedin Airport Musselburgh, Dunedin West Arm, Lake Manapouri Borland Saddle Queenstown Queenstown Aerodrome Mid Dome Cromwell Moa Creek Earnscleugh Alexandra Roxburgh (Power Station) Moa Flat Mahinerangi Dam Tapanui Rankleburn Forest Taieri Mouth Otautau Winton Gore Hokonui Forest Woodlands Invercargill Aerodrome Tiwai Point, Bluff Milton Finegand, Balclutha Finegand, Balclutha Nugget Point Rarotonga Airport, Cook Is. Raoul Island, Kermadec Is. Waitangi, Chatham Islands	24 18 1 2 30 180 180 991 329 352 387 213 427 152 141 110 410 396 2255 15 55 44 72 46 47 0 5 18 6 131 5 5 5 5 6 131 6 131 6 131 6 131 6 131 6 131 131	12.0 11.7 12.0 11.3 11.5 7.1 5.5 9.0 11.3 11.2 12.0 9.8 9.7 11.6 11.6 11.4 11.4 11.4 11.4 11.4 11.4	1.6 1.7 0.8 4.8 3.7 0.2 -0.8 0.0 1.5 1.6 0.7 2.6 1.1 2.4 1.9 2.5 1.3 0.7 4.8 19.1 14.9	6.87 6.41 7.66 8.16 6.87 5.22 6.88 7.19 6.67 7.66 7.66 7.66 7.66 7.66 7.66 7.6	+0.5 +0.5 0.0 +0.8 +0.3 +0.4 +0.5 +0.9 +0.6 +0.7 +0.1 +0.7 +0.1 +0.7 +0.4 +0.8 +0.8 +0.8 +0.8 +0.5 +0.9 +0.5	17.2 16.0 15.9 16.5 9.5 14.8 15.5 17.1 17.2 18.5 18.5 15.6 15.2 16.5 14.5 17.8 17.8 17.8 17.8 17.8 17.0 19.0 17.0 19.0 14.1 27.4 21.4 21.4 21.4 21.8	25 25 26 12 26 26 25 25 25 25 25 25 25 25 25 25 25 25 25	-5.3 -3.6 -0.8 -0.5 -2.5 -4.6 -3.3 -6.5 -7.1 -6.6 -3.8 -4.5 -2.0 -4.5 -4.0 -4.8 -4.2 -4.0 -4.8 -4.2 -4.0 -4.8 -4.2 -1.0	2 3 20 22 33 3 2 2 3 3 4 4 4 3 4 4 4 4 4 4 4 4	79 55 55 641 319 56 37 43 36 29 21 20 36 82 40 58 53 44 52 58 54 37 59 144 275 59	12 13 14 12 14 24 13 14 11 11 9 9 12 12 16 15 16 11 18 18 18 18 18 18 19 19 19 19 19 19 19 19 19 19 19 19 19	+31 +4 +12 +8 +11 +35 -8 -13 +16 -16 +10 -16 +2 -23 -18 -31 -24 -24 -24 -31 -31	19 10 12 15 50 15 13 14 8 16 6 4 4 8 33 13 13 13 13 11 10 8 8 8 10 11 11 8 11 11 11 11 11 11 11 11 11 11 1	23 16 5 23 28 25 31 31 16 5 26 26 31 31 31 31 31 31 31 31 31 31 31 31 31	98 93 115 117 142 120 155 171 86
Campbell Island Scott Base, Antarctica	15 18	8.2	3.3	5.8 -30.4	+0.9 +0.6	10.0 -9.0	26 24	$\begin{array}{c} 4.4 \\ -47.0 \end{array}$	2 19	122	24	••	36		29
					LATE	RETU	RNS								
Te Aroha, July 1971 Tarawera Forest, July 1971	12 61	14.5 14.4	2.0	8.3	-0.8 -0.3	19.0 17.8	27 27	$\begin{bmatrix} -1.7 \\ -4.0 \end{bmatrix}$	14 24	272 125	15	97 66	158	26	
Whatawhata, July 1971 Ruatoria, July 1971 Makaretu, July 1971 Ski Basin, Craigieburn, July	104 61 335	13.7 14.0 10.9	4.1 5.7 3.4	9.0 9.9 7.2	$+0.6 \\ +0.7 \\ \cdots$	16.5 19.4 16.2	27 27 27 27	$ \begin{array}{c c} -2.5 \\ 0.4 \\ 0.5 \end{array} $	17 7 19	129 314 98	10 20 25	$-36 \\ +121 \\ \cdots$	62 93 22	2 2 2 2	i51
1971 Craigieburn Forest, July 1971	1,555 915	0.5 6.1	$\begin{bmatrix} -4.3 \\ -2.5 \end{bmatrix}$	-1.9 1.8	$^{+0.2}_{+0.7}$	5.0 10.4	25 25	$^{-10.2}_{-7.3}$	8 9	32	15	-80	9	28	• •
Camp Stream, Craigicburn, July 1971 Lake Pukaki No. 2, July 1971 Adair, July 1971 Lower Retaruke, June 1971	1,433 556 85 223	1.0 4.7 9.3 14.6	$ \begin{array}{c c} -2.3 \\ -1.9 \\ 1.7 \\ 5.7 \end{array} $	$ \begin{array}{c c} -0.7 \\ 1.4 \\ 5.5 \\ 10.1 \end{array} $	0.0 -0.2	5.6 10.9 15.1 19.5	25 28 28 4	-7.7 -8.5 -2.1 -1.8	8 11 10 21	 8 16 131	6 11 23	 -27 	6 6 25	 4 14 28	

The "normal" refers to the present site of the instruments. The standard periods for normals are: for temperature 1931-60, for rainfall 1921-50, and for sunshine 1935-60. No normals are available for stations with only short records.

Notes on the Weather for August 1971

General—August was unusually mild, especially in the North Island, and it brought to an end a particularly mild winter. Farming conditions varied considerably. In some northern and eastern districts of the North Island and also in Otago the ground was too wet. In Northland, in eastern districts of the North Island, and about Cook Strait serious losses of new-born lambs were reported, some in a violent south-easterly gale with rain affecting especially the Cook Strait area from the 16th to the 18th.

This was another month with an unusually high frequency of

north-easterly flow.

Rainfall—In the North Island rainfall was mainly up to 50 percent above normal, but in the South Island it was predominantly 25 percent below normal.

Central Canterbury with Banks Peninsula had less than half the normal value. On the other hand, several areas had more than double the normal value, namely Northland, from south of Kaitaia to Whangarei, the East Coast as far south as Tokomaru Bay, and the area about Cook Strait.

Northland received its heaviest rain on the 2nd and 3rd, some stations recording as much as 200 mm over these 2 days; and flooding was reported. The East Coast's heavy rain was on the 7th. The area about Cook Strait received its heavy rain on the 16th and 17th; Wellington Airport was drenched by 131 mm in 12 hours.

Temperatures—Temperatures were well above normal, except in Fjordland and on the Southland and Otago coast, where they were close to normal. Departures averaged 1½°C over the North Island, reaching more than 2°c around Rotorua and Waiouru. Over most

^{*}The sunshine recorder is not located at the station but is in the near vicinity.